

ORIGINAL ARTICLE

Cytochrome P450-mediated metabolism of *N*-(2-methoxyphenyl)-hydroxylamine, a human metabolite of the environmental pollutants and carcinogens *o*-anisidine and *o*-nitroanisole

Karel NAIMAN¹, Helena DRAČÍNSKÁ¹, Martin DRAČÍNSKÝ², Markéta MARTÍNKOVÁ¹, Václav MARTÍNEK¹, Petr HODEK¹, Martin ŠTÍCHA³, Eva FREI⁴, Marie STIBOROVÁ¹

¹ Department of Biochemistry, Faculty of Science, Charles University, Prague, Czech Republic

² Institut of Organic Chemistry and Biochemistry, v.v.i., Academy of Sciences, Prague, Czech Republic

³ Department of Organic Chemistry, Faculty of Science, Charles University, Prague, Czech Republic

⁴ Division of Molecular Toxicology, German Cancer Research Center, Heidelberg, Germany

ITX013408A03 • Received: 28 November 2008 • Revised: 5 December 2008 • Accepted: 7 December 2008 • Electronic Publication: November 2010

ABSTRACT

N-(2-methoxyphenyl)hydroxylamine is a human metabolite of the industrial and environmental pollutants and bladder carcinogens 2-methoxyaniline (*o*-anisidine) and 2-methoxynitrobenzene (*o*-nitroanisole). Here, we investigated the ability of hepatic microsomes from rat and rabbit to metabolize this reactive compound. We found that *N*-(2-methoxyphenyl)hydroxylamine is metabolized by microsomes of both species mainly to *o*-aminophenol and a parent carcinogen, *o*-anisidine, whereas 2-methoxynitrosobenzene (*o*-nitrosoanisole) is formed as a minor metabolite. Another *N*-(2-methoxyphenyl)hydroxylamine metabolite, the exact structure of which has not been identified as yet, was generated by hepatic microsomes of rabbits, but its formation by those of rats was negligible. To evaluate the role of rat hepatic microsomal cytochromes P450 (CYP) in *N*-(2-methoxyphenyl)hydroxylamine metabolism, we investigated the modulation of its metabolism by specific inducers of these enzymes. The results of this study show that rat hepatic CYPs of a 1A subfamily and, to a lesser extent those of a 2B subfamily, catalyze *N*-(2-methoxyphenyl)hydroxylamine conversion to form both its reductive metabolite, *o*-anisidine, and *o*-aminophenol. CYP2E1 is the most efficient enzyme catalyzing conversion of *N*-(2-methoxyphenyl)hydroxylamine to *o*-aminophenol.

KEY WORDS: *o*-anisidine; *N*-(2-methoxyphenyl)hydroxylamine; metabolism; oxidation; cytochrome P450

Introduction

2-Methoxyaniline (*o*-anisidine) is a potent carcinogen, causing tumors of the urinary bladder in both genders of F344 rats and B6C3F1 mice (NTP, 1978; IARC, 1982). The International Agency for Research on Cancer (IARC) has classified *o*-anisidine as a group 2B carcinogen (IARC, 1982), which is possibly carcinogenic to humans. Besides its carcinogenicity it exhibits other toxic effects, including hematological changes, anemia and nephrotoxicity (NTP,

1978; IARC, 1982). *o*-Anisidine is used as an intermediate in the manufacturing of a number of azo and naphthol pigments and dyes, which are used for printing (90%) and for paper (3%) and textile (7%) dyeing (NTP, 1978; Garner *et al.*, 1984). Such a wide use of this aromatic amine could result in occupational exposure. Furthermore, it may be released from textiles and leather goods colored with these azo dyes and a large part of the population may be exposed. This carcinogen is also a constituent of cigarette smoke (IARC, 1982; Stabbert *et al.*, 2003). This strongly suggests that *o*-anisidine ranks not only among occupational pollutants produced in the manufacturing of chemicals, but also among environmental pollutants; it can be assumed that human exposure is widespread. Indeed, *o*-anisidine was found in human urine samples in the general population, in concentrations of 0.22 µg/l (median) (Weiss and Angerer, 2003). In addition, hemoglobin adducts of *o*-anisidine were detected in blood samples of persons living in urban or rural

Correspondence address:

Prof. Marie Stiborová, DrSc.

Department of Biochemistry, Faculty of Science, Charles University, Prague, Albertov 2030, 128 40 Prague 2, Czech Republic

E-MAIL: stiborov@natur.cuni.cz

areas of Germany (Falter *et al.*, 1994; Branner *et al.*, 1998; Richter *et al.*, 2001). The adducts as well as *o*-anisidine in urine might originate not only from the sources mentioned above, but also from a possible *o*-anisidine precursor, 2-methoxynitrobenzene (*o*-nitroanisole). This chemical was released into the environment in the course of an accident in a German chemical plant, causing subsequently local and regional contamination (Falter *et al.*, 1994; Hauthal, 1993; Traupe *et al.*, 1997). *o*-Nitroanisole exhibits strong carcinogenic activity, causing neoplastic transformation in the urinary bladder, and to a lesser extent, in the spleen, liver and kidneys in rodents (NTP 1993). It is also a toxic compound, causing anemia. The anemia is characterized by increased levels of methemoglobin and accelerated destruction of erythrocytes (NTP 1993).

Recently, we have found that *o*-anisidine is oxidatively activated by peroxidase and cytochrome P450 (CYP) to species binding to DNA *in vitro* (Stiborová *et al.*, 2001; 2002; 2005; Rýdlová *et al.*, 2005; Naiman *et al.*, 2008). We also demonstrated that *o*-anisidine forms DNA adducts *in vivo*. The same adducts as found in DNA incubated with *o*-anisidine and human microsomes *in vitro* were detected in urinary bladder, the target organ, and to a lesser extent, in liver, kidney and spleen of rats treated with *o*-anisidine (Stiborová *et al.*, 2005). The *o*-anisidine-derived DNA adducts were identified as deoxyguanosine adducts formed from a metabolite of *o*-anisidine, *N*-(2-methoxyphenyl)hydroxylamine, which is generated by oxidation of *o*-anisidine with human, rabbit and rat hepatic microsomes (Stiborová *et al.*, 2005; Rýdlová *et al.*, 2005; Naiman *et al.*, 2008). The same deoxyguanosine adducts were also detected in DNA of the urinary bladder, kidney, liver and spleen of rats treated with *o*-nitroanisole (Stiborová *et al.*, 2004), an oxidized counterpart of *o*-anisidine, and in DNA incubated with *o*-nitroanisole *in vitro* with human and rat hepatic cytosolic enzymes and xanthine oxidase (Stiborová *et al.*, 1998; 2004). These enzymatic systems were found to produce *N*-(2-methoxyphenyl)hydroxylamine after *o*-nitroanisole reduction (Mikšanová *et al.*, 2004). The data indicate that formation of *N*-(2-methoxyphenyl)hydroxylamine, the reactive metabolite of both carcinogens, is critical for generation of DNA lesions in target organs. Therefore, it is clear that *N*-(2-methoxyphenyl)hydroxylamine formation and its further conversion, as well as the enzymes participating in such processes, play a key role in carcinogenic effects of both carcinogens.

Recently, we have found that *o*-anisidine is oxidized by human, rat and rabbit hepatic microsomes not only to *N*-(2-methoxyphenyl)hydroxylamine, but that this compound is a subject of complex redox cycling reactions, forming also *o*-aminophenol, *o*-nitrosoanisole and one additional metabolite, the exact structure of which has not been identified as yet (Stiborová *et al.*, 2005; Naiman *et al.*, 2008). *N*-(2-methoxyphenyl)hydroxylamine might also be a subject of complex reactions, and its fate is dependent on the environment, in which it occurs. It can be further metabolized to *o*-aminophenol, *o*-nitrosoanisole and parental *o*-anisidine (Naiman *et al.*, 2008), or when nucleophiles such as DNA or proteins are present in the cell, form the adducts (Stiborová *et al.*, 2005) (Figure 1).

The results of our former studies (Stiborová *et al.*, 2005; Naiman *et al.*, 2008) show a similarity among hepatic microsomal CYP systems metabolizing *o*-anisidine in humans and two animal models, rats and rabbits, which are the species, in which this agent is carcinogenic or toxic (NTP, 1978; IARC, 1982). These findings indicate that both experimental animal species might serve as suitable models to mimic the fate of this carcinogen in human. However, whereas formation of DNA adducts by *N*-(2-methoxyphenyl)hydroxylamine, generated in reactions catalyzed with human and rat hepatic microsomes, has been clearly proven (Stiborová *et al.*, 2005), conversion of this metabolite to further products has been investigated using only rabbit hepatic microsomes (Naiman *et al.*, 2008). Therefore, the aim of the present study was to evaluate the efficiency of *N*-(2-methoxyphenyl)hydroxylamine metabolism by hepatic microsomes from rats.

Materials and methods

Chemicals

Chemicals were obtained from the following sources: β -naphthoflavone (β -NF), NADP⁺, NADPH, glucose 6-phosphate and bicinehoninic acid (2,2'-biquinoline-4,4'-dicarboxylic acid) from Sigma Chemical Co. (St. Louis, MO, USA); *o*-anisidine, *o*-aminophenol (>99% based on HPLC) from Fluka Chemie AG (Buchs, Switzerland) and glucose 6-phosphate dehydrogenase from Serva (Heidelberg, Germany). All these and other chemicals were of analytical purity or better. 2-Methoxynitrosobenzene was synthesized in analogy to the synthesis described earlier (Seidenfaden, 1971) by oxidation of *N*-(2-methoxyphenyl)hydroxylamine with potassium dichromate in water and identified by ¹H NMR recorded on a 400 MHz instrument in CDCl₃ (referenced to TMS): 7.68 (1H, m), 7.36 (1H, m), 6.86 (1H, m), 6.30 (1H, m) and 4.28 (3H, s). *N*-(2-methoxyphenyl)hydroxylamine was synthesized by the procedure similar to that described earlier (Balaban *et al.*, 1998). *N*-(2-Methoxyphenyl)hydroxylamine authenticity was confirmed by electrospray mass and CID spectra and high field proton NMR spectroscopy. The positive-ion electrospray mass-spectrum exhibited the protonated molecule at *m/z* 140.1, while the CID of its ion fragments at *m/z* 125.2, 108.1 and 109.1. The ¹H-NMR spectra were recorded at 400 MHz in dimethyl sulfoxide-*d*₆. The central line of dimethyl sulfoxide at 2.500 ppm was used as reference line. The spectra showed the presence of the following protons: 8.28 (1H, d, J = 2.3 Hz, exchanged with CD₃OD), 7.64 (1H, d, J = 1.5 Hz, exchanged with CD₃OD), 7.01 (1H, m, Σ J = 9.6 Hz), 6.84 (2H, m, Σ J = 15.0 Hz), 6.75 (1H, m, Σ J = 16.9 Hz), 3.75 (3H, s). *o*-Aminophenol authenticity was confirmed by ESI mass spectra and ¹H and ¹³C NMR spectroscopy. The positive-ion mass-spectrum exhibited the protonated molecule at *m/z* 110.4, while the negative-ion mass-spectrum the molecule at *m/z* 108.4. The NMR spectra were recorded on a Bruker Avance II-500 instrument (499.8 MHz for ¹H and 125.7 MHz for ¹³C) in dimethyl sulfoxide-*d*₆ and referenced to the solvent signal (δ 2.50 and 39.70, respectively). ¹H NMR spectrum showed the presence of the following signals: 8.96 (1H, bs), 6.65 (1H, m), 6.59 (1H, m), 6.54 (1H, m), 6.40 (1H,

m) and 4.46 (2H, bs). ^{13}C NMR spectrum showed the presence of six signals of aromatic carbons: 144.26 (s), 136.78 (s), 119.80 (d), 116.75 (d), 114.73 (d) and 114.64 (d).


Animal experiments, preparation of microsomes and assays

The study was conducted in accordance with the Regulations for the Care and Use of Laboratory Animals (311/1997, Ministry of Agriculture, Czech Republic), which complies with Declaration of Helsinki. Microsomes from livers of ten untreated rats and three rabbits were prepared by the procedure described previously (Stiborová *et al.*, 1988). Microsomes from the livers of ten male Wistar rats pretreated with β -NF (Stiborová *et al.*, 1988) were isolated as described (Stiborová *et al.*, 1988; 1995), those pretreated with phenobarbital (PB) as reported by Hodek *et al.* (1988), and those pretreated with ethanol were isolated using a procedure described by Yang *et al.* (1985). Protein concentrations in the microsomal fractions were assessed using the bicinchoninic acid protein assay with bovine serum albumin as a standard (Wiechelman *et al.*, 1988). The concentration of CYP was estimated according to Omura and Sato (1964), by measuring the absorption of the complex of reduced CYP with carbon monoxide. Rat and rabbit liver microsomes contained 0.6 and 1.8 nmol CYP/mg protein, respectively. Hepatic microsomes of rats induced with β -NF, PB, and ethanol contained 1.3, 2.7 and 1.8 nmol CYP/mg protein, respectively. The activity of NADPH:CYP reductase in rat hepatic microsomes was measured according to Sottocasa *et al.* (1967) using cytochrome c as substrate (i.e., as NADPH:cytochrome c reductase). NADPH:CYP reductase activities in hepatic microsomes of control (uninduced) rats and those induced with β -NF, PB and ethanol were 0.210, 0.199, 0.325 and 0.201 $\mu\text{mol}/\text{min}/\text{mg}$ protein, respectively.

Incubations

Incubation mixtures used for study of the *o*-anisidine metabolism contained the following concentrations in a

final volume of 100 μl : 100 mM sodium phosphate buffer (pH 7.4), 1 mM NADP⁺, 10 mM D-glucose 6-phosphate, 1 U/ml D-glucose 6-phosphate dehydrogenase (NADPH-generation system), a rat or rabbit hepatic microsomal fraction containing 0.04–1.0 nmol CYP, and 0.1–2.0 mM *o*-anisidine dissolved in 1.0 μl methanol. The reaction was initiated by adding the substrate. To study metabolism of *N*-(2-methoxyphenyl)hydroxylamine, incubation mixtures contained the following concentrations in a final volume of 100 μl : 100 mM sodium phosphate buffer (pH 7.4), 1 mM NADP⁺, 10 mM D-glucose 6-phosphate, 1 U/ml D-glucose 6-phosphate dehydrogenase (NADPH-generation system), a rat and rabbit hepatic microsomal fraction containing 0.04–1.0 nmol CYP, and 0.1–1.0 mM *N*-(2-methoxyphenyl)hydroxylamine dissolved in 1.0 μl distilled water. The reaction was initiated by adding the substrate. After incubation in open glass tubes (37°C, 30 min), the reactions were terminated by adding 100 μl of methanol and centrifuged at 5,000 g for 5 min. Metabolism of *o*-anisidine and *N*-(2-methoxyphenyl)hydroxylamine with rat and rabbit CYP enzymatic systems was linear until 40 min. The supernatants were collected and 20 μl aliquots applied onto a high-performance liquid chromatography (HPLC) column, where metabolites of *o*-anisidine were separated. The HPLC was performed on a C-18 reversed-phase column (250 \times 4.6 mm, 5 μm , Nucleosil 100-5, Macherey-Nagel, Duren, Germany). Metabolites were eluted with 18% methanol, 82% 7.18 μM aqueous ammonia, pH 8.0, (v/v) at a flow rate of 0.6 ml/min and monitored at 254 nm. *o*-Anisidine and *N*-(2-methoxyphenyl)hydroxylamine metabolites were analyzed by mass spectrometry and by comparing their chromatographic properties on HPLC with those of synthetic standards. 2-Methoxynitrosobenzene, *o*-aminophenol, *N*-(2-methoxyphenyl)hydroxylamine and 2-methoxynitrosobenzene standards, were eluted at retention times (r.t.) of 8.8, 11.3, 19.7 and 57.5 min, respectively (Stiborová *et al.*, 2005; Naiman *et al.*, 2008).



Results

Metabolism of *N*-(2-methoxyphenyl)hydroxylamine by rat and rabbit hepatic microsomes

When a parental compound from which *N*-(2-methoxyphenyl)hydroxylamine is generated, *o*-anisidine, was incubated with rat and rabbit hepatic microsomes in the presence of NADPH, two metabolites, *o*-aminophenol (r.t. of 11.3 min) and *N*-(2-methoxyphenyl)hydroxylamine (r.t. of 19.7 min), as well as an additional product peak of metabolite 1 (M1, r.t. of 7.0 min), which structure has not been identified as yet, were separated by HPLC (see peaks in Figure 2A for the profile obtained with rabbit microsomes) (Naiman *et al.*, 2008).

The results of experiments with *N*-(2-methoxyphenyl)hydroxylamine and microsomes demonstrated that metabolites M1 and *o*-aminophenol are also formed from this *o*-anisidine metabolite by rabbit hepatic microsomes (Figure 2B), while formation of the metabolite M1 from this compound by hepatic microsomes of rats was negligible (Figure 2C). When *N*-(2-methoxyphenyl)hydroxylamine was incubated without hepatic microsomal enzymes or without NADPH, metabolite M1 and *o*-aminophenol peaks were also detectable by HPLC, but only under acidic conditions (pH 4.5 for 60 min). At pH 7.4, used for microsomal incubations, their spontaneous formation was negligible. This finding indicates that conversion of *N*-(2-methoxyphenyl)hydroxylamine in microsomes is mediated by enzymatic reactions. During metabolism of *N*-(2-methoxyphenyl)hydroxylamine by hepatic microsomes of both species, a shoulder at 8.8 min was also detectable (Figure 2B,C), suggesting formation of *o*-nitrosoanisole (2-methoxynitrosobenzene) (r.t. of 8.8 min, Figure 2F). Moreover, during the incubations of *N*-(2-methoxyphenyl)hydroxylamine with rat and rabbit microsomes and NADPH, an additional product peak was detected by HPLC, being identified to be the parental compound, *o*-anisidine (r.t. of 28.6 min) (Figure 2B,C).

Involvement of CYP enzymes in *N*-(2-methoxyphenyl)hydroxylamine metabolism in rat hepatic microsomes

In order to evaluate the participation of individual rat hepatic microsomal CYPs in *N*-(2-methoxyphenyl)hydroxylamine metabolism, the induction of individual CYP enzymes was performed with this animal model. Microsomes isolated from livers of uninduced rats and rats pre-treated with β -NF (enriched with CYP1A1/2), PB (enriched with CYP2B1/2) and ethanol (enriched with CYP2E1) were used (Table 1).

Except of the enzymes present in microsomes of livers of rats pre-treated with ethanol, the enzymes of all other microsomes tested in this study formed from *N*-(2-methoxyphenyl)hydroxylamine predominantly its reductive metabolite, *o*-anisidine. More than 2-fold higher levels of *o*-anisidine than *o*-aminophenol were determined when *N*-(2-methoxyphenyl)hydroxylamine was incubated with these microsomes and NADPH (Table 1). Levels of the *N*-(2-methoxyphenyl)hydroxylamine reductive metabolite, *o*-anisidine, did not correspond to activities of NADPH:CYP reductase in individual microsomes. Namely, activities of this enzyme in hepatic microsomes of control (uninduced)

rats and those induced with β -NF, PB and ethanol used in the experiments are similar, being 0.210, 0.199, 0.325 and 0.201 μ mol/min/mg protein, respectively. While incubations of *N*-(2-methoxyphenyl)hydroxylamine with hepatic microsomes of rats pre-treated with β -NF led to a 2.4- and 1.9-fold increase ($P < 0.05$) in *o*-aminophenol and *o*-anisidine formation, respectively (Table 1), another inducer of CYP enzymes, PB, had much less effect. Even though a 1.4- and 1.2-fold increase in *o*-aminophenol and *o*-anisidine formation, respectively, was mediated by treating rats with this CYP inducer (Table 1), this increase was statistically insignificant. Ethanol, an inducer of CYP2E1, stimulated production of *o*-aminophenol from *N*-(2-methoxyphenyl)hydroxylamine, by 3-fold ($P < 0.05$), whereas decreased levels of the *N*-(2-methoxyphenyl)hydroxylamine reduction metabolite, *o*-anisidine. These results indicate that rat hepatic CYP2E1, CYPs of a 1A subfamily and, to a lesser extent, those of a 2B subfamily, are capable of metabolizing *N*-(2-methoxyphenyl)hydroxylamine in rat livers.

It should be noted that the results of experiments with inducers should be carefully interpreted, because these inducers are not absolutely specific for individual CYPs. Therefore, to confirm the role of these CYPs in *N*-(2-methoxyphenyl)hydroxylamine metabolism, additional experimental approaches such as selective inhibition of CYPs and utilization of the purified CYP reconstituted with NADPH: CYP reductase are planned to be employed in further studies.

Discussion

The results of this study show that rat and rabbit hepatic microsomes can metabolize *N*-(2-methoxyphenyl)hydroxylamine, a reactive metabolite of carcinogenic *o*-anisidine and *o*-nitrosoanisole. This compound is responsible for genotoxic effects of both carcinogens, because it is easily decomposed to the nitrenium/carbenium ion forming DNA adducts (Figure 1) (Stiborová *et al.*, 2004; 2005; Naiman *et al.*, 2008). The results demonstrate that *N*-(2-methoxyphenyl)hydroxylamine is also further metabolized to *o*-aminophenol, *o*-nitrosoanisole and the parent compound, *o*-anisidine. The

Table 1. Metabolism of *N*-(2-methoxyphenyl)hydroxylamine in rat hepatic microsomes induced with different agents

Hepatic microsomes	<i>N</i> -(2-methoxyphenyl)hydroxylamine metabolites ^a	
	<i>o</i> -aminophenol	<i>o</i> -anisidine
from rats pretreated with ^b		
None - control microsomes	2.0 \pm 1.0	5.8 \pm 0.7
β -naphthoflavone (CYP1A1/2)	4.8 \pm 1.7	11.1 \pm 3.3
Phenobarbital (CYP2B1/2)	2.8 \pm 2.5	7.0 \pm 2.0
Ethanol (CYP2E1)	6.1 \pm 3.1	3.5 \pm 0.7

^a The numbers are the peak area/min/nmol CYP for each metabolite; averages \pm S.E.M of three determinations in separate experiments.

^b Isoforms of CYP induced are shown in brackets.




Figure 2. HPLC elution profiles of metabolites of 1 mM *o*-anisidine incubated with rabbit microsomes (A), of 1.0 mM N-(2-methoxyphenyl)hydroxylamine incubated with rabbit (B) and rat (C) hepatic microsomes. (D) synthetic N-(2-methoxyphenyl)hydroxylamine and *o*-anisidine. (E) *o*-aminophenol. (F) *o*-nitrosoanisole. For incubation conditions see *Materials and methods*. Peaks eluting between 2.0 and 5.5 min, solvent front, NADPH and protein components of microsomes and NADPH-generation system.

formed *o*-anisidine may be *O*-demethylated again to *o*-aminophenol (Figure 1). The question whether *o*-aminophenol is also formed from *N*-(2-methoxyphenyl)hydroxylamine by its *O*-demethylation to *N*-(2-hydroxyphenyl)hydroxylamine, which is subsequently reduced to *o*-aminophenol (Figure 1), remains to be answered. No metabolites formed by this reaction were observed.

Recently, redox cycling reactions similar to those we found with *N*-(2-methoxyphenyl)hydroxylamine were observed by Kim *et al.* (2004), who studied metabolism of several aromatic and heterocyclic amines by a CYP1A2/NADPH:CYP reductase enzymatic system. They reported that the CYP system catalyzes oxidation of the *N*-hydroxylated intermediate formed from the carcinogenic heterocyclic amine 2-amino-3-methylimidazo[4,5-*f*]quinoline (IQ), to a nitrosoderivative. They demonstrated that NADPH:CYP reductase can catalyze the reduction of the IQ oxidation products, *N*-nitroso-IQ and *N*-hydroxyl-IQ, to *N*-hydroxyl-IQ and the parent amine, IQ (Kim *et al.*, 2004). *N*-hydroxylation products of two other aromatic amines investigated by Kim *et al.* (2004), 2-aminofluorene and 4-aminobiphenyl, are, however, reduced non-enzymatically, by NADPH. We have not determined whether reduction of *N*-(2-methoxyphenyl)hydroxylamine to *o*-anisidine requires catalysis by NADPH:CYP reductase or occurs non-enzymatically, or is mediated by other enzymes. However, preliminary experiments performed in our laboratory suggest that although NADPH:CYP reductase might partially participate in *N*-(2-methoxyphenyl)hydroxylamine reduction to *o*-anisidine, CYP enzymes present in hepatic microsomes are more effective in this process. (Naiman *et al.*, unpublished results). Indeed, the results of the present study demonstrate that rat hepatic microsomes reduce *N*-(2-methoxyphenyl)hydroxylamine to *o*-anisidine independently on activities of NADPH:CYP reductase. The present study shed also some light on the role of specific microsomal CYP enzymes in metabolism of *N*-(2-methoxyphenyl)hydroxylamine. The CYP1A enzymes seem to be most efficient in *N*-(2-methoxyphenyl)hydroxylamine reduction. Nevertheless, the question whether CYP2E1 is also effective in reducing *N*-(2-methoxyphenyl)hydroxylamine, remains to be answered. Among hepatic microsomes tested in this study, those of rats treated with ethanol (enriched with CYP2E1) produced the lowest levels of *o*-anisidine from *N*-(2-methoxyphenyl)hydroxylamine. In contrast, *o*-aminophenol was the major metabolite in microsomes enriched with CYP2E1, generated at the highest levels among the microsomes employed in the experiments. These findings suggest that CYP2E1 is not effective in *N*-(2-methoxyphenyl)hydroxylamine reduction. Nevertheless, because CYP2E1 is the most effective CYP enzyme oxidizing *o*-anisidine to *o*-aminophenol (Stiborová *et al.*, 2005; Naiman *et al.*, 2008), one can speculate that this enzyme efficiently utilizes *o*-anisidine, which could be generated in this microsomal system, oxidizing it to *o*-aminophenol. Therefore, the participation of CYP2E1 in *N*-(2-methoxyphenyl)hydroxylamine reduction to *o*-anisidine cannot be excluded and awaits further investigation. The study utilizing purified CYP2E1 reconstituted

with NADPH:CYP reductase is planned to be performed to explain this question.

While the formation of *N*-(2-methoxyphenyl)hydroxylamine was clearly identified to be the activation pathway of *o*-anisidine and *o*-nitroanisole metabolism (Stiborová *et al.*, 2004; 2005), biological significance of formation of *o*-aminophenol for detoxication/activation metabolism awaits further investigation. *o*-Aminophenol might be considered to be mutagenic, because it induces sister chromatid exchanges in a dose-dependent manner in cultured human lymphocytes *in vitro* and in Chinese hamster bone marrow cells *in vivo* (Kirchner and Bayer, 1992). In addition, Brennan and Schiestl (1997) reported that *o*-aminophenol is positive in the deletion recombination assay in *Saccharomyces cerevisiae*. Even though *o*-aminophenol has not been found to form covalent DNA adducts, it was demonstrated in *in-vitro* experiments to cause DNA damage, forming 8-oxy-7,8-dihydro-2'-deoxyguanosine in the presence of metal ions such as Cu(II) (Okhuma and Kawanishi, 2001). Hence, due to such processes, *o*-aminophenol may contribute to initiation of the *o*-anisidine- and/or *o*-nitroanisole-mediated carcinogenesis in the urinary bladder, and in a tumor development induced by other bladder carcinogenic aromatic amines, which produce this compound as one of the metabolites (Brennan and Schiestl, 1997). Furthermore, *O*-demethylation reactions produce formaldehyde (Figure 1), which is known to modify DNA, generating several products including hydroxymethyl adducts and cross-links (Beland *et al.*, 1984; Huang and Hopkins, 1993; Cheng *et al.*, 2003). Formaldehyde is mutagenic in a variety of different test systems and carcinogenic in laboratory animals (IARC, 2006) and has been described as "carcinogenic to human" by the IARC and "reasonably anticipated to be a human carcinogen" by the U.S. Department of Health and Human Services (2004). Therefore, on the one hand, it is plausible that formaldehyde-DNA adducts could also play a role in carcinogenesis by *o*-anisidine. On the other hand, however, formaldehyde produced in the cell is also detoxified by conjugation to glutathione and oxidized (Dharaeswar and Stella, 2008); therefore, it is not likely a strong contributor to carcinogenicity caused by *o*-anisidine.

It should be noted that tumor development in a specific organ is influenced by promotional pressures on initiated cells in target organs and not only by the levels of DNA adducts formed by the compounds like *o*-anisidine and *o*-nitroanisole. It is known that radicals formed from several carcinogens producing oxidative DNA damage, such as 8-hydroxy-2'-deoxyguanosine, are important not only in initiation, but also in promotion phases of carcinogenesis (Imaoka *et al.*, 2004). Therefore, their formation from *o*-aminophenol may be one of the factors contributing to tumor promotion in *o*-anisidine- and/or *o*-nitroanisole-mediated carcinogenesis. In addition, *o*-anisidine is oxidized by several peroxidases, which are expressed in target organs (e.g. COX), to form radicals besides DNA adducts (Brennan, and Schiestl, 1999; Sasaki *et al.*, 1998). Hence, the production of such free radicals in or near the target cells may be another factor important in the promotional process in *o*-anisidine-mediated tumor development. However, the exact functions

of such and/or other promotional pressures caused both by *o*-anisidine and *o*-aminophenol in an *o*-anisidine-mediated tumorigenesis remain to be resolved.

Acknowledgement

The work is supported by Grant Agency of Charles University (grant 7418/2007), the Grant Agency of the Czech Republic (grants 203/06/0329 and 303/06/0928) and the Ministry of Education of the Czech Republic (grants MSM0021620808 and 1M0505).

REFERENCES

- Balaban AT, Garfield RE, Lesko MJ, Seitz WA. (1998) Synthesis and spectral data of some new N-nitroso-N-phenylhydroxylamine (cupferron) derivatives. *Org Prep Proced Int* **30**: 439–446.
- Beland FA, Fullerton NF, Heflich RH. (1984) Rapid isolation, hydrolysis and chromatography of formaldehyde-modified DNA. *J Chromatogr* **308**: 121–131.
- Branner B, Kutzer C, Zwicknnpflug W, Scherer G, Heller W-D, Richter E. (1998) Haemoglobin adducts from aromatic amines and tobacco-specific nitrosamines in pregnant smoking and non-smoking women. *Biomarkers* **3**: 35–47.
- Brennan RJ, Schiestl RH. (1997). Aniline and its metabolite generate free radicals in yeast. *Mutagenesis* **12**: 215–220.
- Brennan RJ, Schiestl RH. (1999) The aromatic amine carcinogens *o*-toluidine and *o*-anisidine induce free radicals and intrachromosomal recombination in *Saccharomyces cerevisiae*. *Mutat Res* **430**: 37–45.
- Cheng G, Shi Y, Sturla S, J alas J, McIntee EJ, Villalta PW, Wang M, Hecht SS. (2003) Reactions of formaldehyde plus acetaldehyde with deoxyguanosine and DNA: Formation of cyclic deoxyguanosine adducts and formaldehyde cross-links. *Chem Res Toxicol* **16**: 145–152.
- Dhareshwar SS, Stella VJ. (2008) Your prodrug releases formaldehyde: Should you be concerned? No! *J Pharm Sci* **97**: 4184–4193.
- Falter B, Kutzer C, Richter E. (1994) Biomonitoring of hemoglobin adducts: aromatic amines and tobacco-specific nitrosamines. *Clin Invest* **72**: 364–371.
- Garner RC, Martin CN, Clayson DB (1984) (2nd ed. Searle C, Ed.) Carcinogenic aromatic amines and related compounds. Chemical Carcinogens, ACS Monograph 182, Vol. 1: pp. 175–302, American Chemical Society, Washington, DC.
- Hauthal HG. (1993) Rosenmontag und die Folgen die Störfälle der Hoechst AG. *Nach. Chem Tech Lab* **41**: 440.
- Hodek P, Jansčák P, Anzenbacher P, Burkhard J, Janků J, Vodička L. (1988) Metabolism of diamantane by rat-liver microsomal-cytochromes P-450. *Xenobiotica* **18**: 1109–1118.
- Huang H, Hopkins PB. (1993) DNA interstrand cross-linking by formaldehyde: nucleotide sequence preference and covalent structure of the predominant cross-link formed in synthetic oligonucleotides. *J Am Chem Soc* **115**: 9402–9408.
- Imaoka S, Osada M, Minamiyama Y, Yukimura T, Toyokuni S, Takemura S, Hiroi T, Funae Y. (2004) Role of phenobarbital-inducible cytochrome P450s as a source of active oxygen species in DNA-oxidation. *Cancer Lett* **203**: 117–125.
- IARC (1982) Ortho- and para-anisidine and their hydrochlorides. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans, No. 27, IARC, Lyon.
- IARC (2006) Formaldehyde, 2-butoxyethanol and 1-tert-butoxypropan-2-ol. IARC Monographs on the Evaluation of Carcinogenic Risk to Humans, Vol. 88, pp. 39–325, IARC, Lyon.
- Kim D, Kadlubar FF, Teitel CH, Guengerich FP. (2004) Formation and reduction of aryl and heterocyclic nitroso compounds and significance in the flux of hydroxylamines. *Chem Res Toxicol* **17**: 529–536.
- Kirchner G, Bayer U. (1992) Genotoxic activity of the aminophenols as evidenced by the induction of sister chromatid exchanges. *Hum Toxicol* **1**: 387–392.
- Mikšanová M, Novák P, Frei E, Stiborová M. (2004) Metabolism of carcinogenic 2-nitroanisole by rat, rabbit, porcine and human hepatic cytosol. *Collect Czech Chem Commun* **69**: 589–602.
- Naiman K, Dračínská H, Martínková M, Šulc M, Dračínský M, Kejíková L, Hodek P, Hudeček J, Liberda J, Schmeiser HH, Frei E, Stiborová M. (2008) Redox cycling in the metabolism of the environmental pollutant and suspected human carcinogen *o*-anisidine by rat and rabbit hepatic microsomes. *Chem Res Toxicol* **21**: 1610–1621.
- NTP (1978) Bioassay of *o*-anisidine hydrochloride for possible carcinogenicity. NTP Technical Report 89, National Institute of Health, US Department of Health and Human Services, Bethesda, MD.
- NTP (1993) Toxicology and Carcinogenesis. Studies of 2-nitroanisole. NTP Technical Report 416, National Institute of Health, US Department of Health and Human Services, Bethesda, MD.
- Okhuma Y, Kawanishi S. (2001) Oxidative DNA damage induced by a metabolite of carcinogenic *o*-anisidine: enhancement of DNA damage and alteration in its sequence specificity by superoxide dismutase. *Arch Biochem Biophys* **389**: 49–56.
- Omura T, Sato R. (1964) The carbon monoxide-binding pigment of liver microsomes. I. Evidence for its hemoprotein nature. *J Biol Chem* **239**: 2370–2378.
- Richter E, Rosler S, Scherer G, Gostomzyk JG, Grubl A, Kamer U, Behrendt H. (2001) Haemoglobin adducts from aromatic amines in children in relation to area of residence and exposure to environmental tobacco smoke. *Int Arch Occup Environ Health* **74**: 421–428.
- Rýdlová H, Mikšanová M, Ryšlavá H, Stiborová M. (2005) Carcinogenic pollutants *o*-nitroanisole and *o*-anisidine are substrates and inducers of cytochromes P450. *Biomed Papers* **49**: 441–447.
- Sasaki YF, Nishidate E, Su YQ, Matsusaka N, Tsuda S, Susa N, Furukawa Y, Ueno S. (1998) Organ-specific genotoxicity of the potent rodent bladder carcinogens *o*-anisidine and *p*-cresidine. *Mutat Res* **412**: 155–160.
- Seidenfaden W. (1971) Methoden der Organischen Chemie (Houben-Weyl), Band X/1, Georg Thieme Verlag, Stuttgart, p. 1058.
- Sottocasa GL, Kuylenstierna I, Ernster I, Bergstrand A. (1967) An electron transport system associated with the outer membrane of liver mitochondria. *J Cell Biol* **32**: 415–438.
- Stabbert R, Schäfer K-H, Biefel C, Rustemeier K. (2003) Analysis of aromatic amines in cigarette smoke. *Rapid Commun Mass Spectrom* **17**: 2125–2132.
- Stiborová M, Asfaw B, Anzenbacher P, Lešetický L, Hodek P. (1988) The first identification of the benzenediazonium ion formation from a non-aminoazo dye, 1-phenylazo-2-hydroxynaphthalene (Sudan I) by microsomes of rat livers. *Cancer Lett* **40**: 319–326.
- Stiborová M, Asfaw B, Frei E, Schmeiser HH, Wiessler M. (1995) Benzenediazonium ion derived from Sudan I forms an 8-(phenylazo)guanine adduct. *Chem Res Toxicol* **8**: 489–498.
- Stiborová M, Mikšanová M, Havlíček V, Schmeiser HH, Frei E. (2002) Mechanism of peroxidase-mediated oxidation of carcinogenic *o*-anisidine and its binding to DNA. *Mutat Res* **500**: 49–66.
- Stiborová M, Mikšanová M, Smrček S, Bieler CA, Breuer A, Klokow KA, Schmeiser HH., Frei E. (2004) Identification of a genotoxic mechanism for 2-nitroanisole carcinogenicity and of its carcinogenic potential for humans. *Carcinogenesis* **25**: 833–840.
- Stiborová M, Mikšanová M, Šulc M, Rýdlová H, Schmeiser HH, Frei E. (2005) Identification of a genotoxic mechanism for the carcinogenicity of the environmental pollutant and suspected human carcinogen *o*-anisidine. *Int J Cancer* **116**: 667–678.
- Stiborová M, Schmeiser HH, Breuer A, Frei E. (2001) Evidence for activation of carcinogenic *o*-anisidine by prostaglandin H synthase: ³²P-postlabelling analysis of DNA adducts formation. *Gen Physiol Biophys* **20**: 267–279.
- Stiborová M, Schmeiser HH, Frei E. (1998) To the mechanism of 2-nitroanisole carcinogenicity: *in vitro* metabolism of 2-nitroanisole mediated by peroxidases and xanthine oxidase. *Collect Czech Chem Commun* **63**: 857–869.
- Traupe H, Menge G, Kandt J, Karmaus W. (1997) Higher frequency of atopic dermatitis and decrease in viral warts among children exposed to chemicals liberated in a chemical accident in Frankfurt, Germany. *Dermatology* **195**: 112–118.
- U.S. Department of Health and Human Services (2004) 11th Report on Carcinogens, U.S. Government Printing Office, Washington, DC.
- Weiss T, Angerer J. (2003) Simultaneous determination of various aromatic amines and metabolites of aromatic nitro compounds in urine for low level exposure using gas chromatography-mass spectrometry. *J Chromatogr B* **778**: 179–192.
- Wiechelmann KJ, Braun RD, Fitzpatrick JD. (1988) Investigation of the bicinchoninic acid protein assay: identification of the groups responsible for color formation. *Anal Biochem* **175**: 231–237.
- Yang CS, Tu YY, Koop DR, Coon MJ. (1985) Metabolism of nitrosamines by purified rabbit liver cytochrome P-450 isozymes. *Cancer Res* **45**: 1140–1145.