

ORIGINAL ARTICLE

Toxicity of perfluorinated carboxylic acids for aquatic organisms

Miloň TICHÝ¹, Radka VALIGUROVÁ², Radomír ČABALA², Rut UZLOVÁ¹, Marián RUCKI¹

¹ Laboratory of Predictive Toxicology, National Institute of Public Health, Praha, Czech Republic

² Charles University in Prague, Faculty of Science, Department of Analytical Chemistry, Praha, Czech Republic

ITX030210A01 • Received: 10 June 2010 • Revised: 20 June 2010 • Accepted: 21 June 2010

ABSTRACT

Toxicity of perfluorinated carboxylic acids with carbon chain C₈ to C₁₂ were tested with oligochaeta *Tubifex tubifex*. Toxicity was evaluated as the exposure time ET₅₀ from onset of damage of the oligochaeta in saturated aqueous solutions. The ET₅₀ fluctuated between 25 and 257 minutes. No statistically significant difference was found among the C₈, C₉ and C₁₂ acids (ET₅₀ between 143 and 257 minutes with large standard deviation). The acids with carbon chain C₁₀ and C₁₁ induced the effect significantly quicker (25 to 47 minutes). No acute toxicity measured in the three-minute test was observed in any case.

KEY WORDS: perfluorinated carboxylic acids; *Tubifex tubifex*; water pollution; effective exposure time; acute toxicity

This paper is dedicated to the memory of Prof. Helena Rašková, MD., DSc.

Introduction

It is impossible to mention all the fluorinated products used nowadays in miscellaneous human activities, whereas naturally occurring ones are more than rare. Moreover, fluorinated derivatives of organic compounds belong to the most stable compounds, badly degradable, easily distributed in various media and matrices over the world. Their unique chemical properties make them important ingredients in various products. The compounds are found as widespread environmental contaminants. The global-scale multispecies mass balance model was used for simulation of the long-term transport of polyfluorinated carboxylic acids containing 8 to 13 carbons (Armitage *et al.*, 2009). Model scenarios estimated their direct emission and indicated that the mass fluxes to the Arctic marine environment associated with oceanic transport were in excess of mass fluxes from indirect sources (atmospheric transport of precursors such as

fluorotelomer alcohols and subsequent degradation to perfluorinated carboxylic acids, PFCAs).

Organic fluorinated acids are present in both arctic and antarctic ice or in equatorial waters. The polyfluoroalkyl acids and, in the past, fluoro-chloro methanes or ethanes belong among the most extended ones. The fluoro-halogen alkanes were used in sprays, their ethers as anesthetics, perfluorooctanoic and perfluorooctane sulphonate acids are widely used today in a lot of products as repellents, paper coatings, coatings for leather, upholstery or carpets, in fire fighting foams, pharmaceuticals, insecticides, *etc.*, (Alloatti *et al.*, 2008, Houde *et al.*, 2006, Kovarova & Svobodova, 2008).

On the other hand, monofluorinated organic compounds are important in clinical disciplines. Thus *e.g.* during the last decades, 2-deoxy-2-fluoro-D-glucose has been used to assess glucose uptake in the normal and diseased heart muscle (Hariharan *et al.*, 1995), fluoro-derivatives of benzimidazole-quinoline were found to be inhibitors of the multitargeted receptor tyrosine kinase and on aspiration to be an anticancer drug (Henk *et al.*, 2005), or fluoro-derivatives of 1,4-benzoquinone were used for studying the participation of ubiquinone in bovine heart mitochondrial respiratory chain, among many other products. The introduction of fluorine atoms into biologically active compounds proved to be

Correspondence address:

Assoc. Prof. Miloň Tichý, DSc.

Laboratory of Predictive Toxicology, National Institute of Public Health,
10042 Praha 10, Czech Republic

E-MAIL: mtichy@szu.cz

beneficial. Several antitumor agents of this type can be named, such as fluorouracil, floxuridin, tegafur, getitinib or camptothecin (Alloatti *et al.*, 2008).

And all these compounds eventually end up in our environment.

Generally, acute toxicity of polyfluorinated compounds is low (Kovarova and Svobodova, 2008), yet there is a lack of toxicological data. There is also a lack of data on their physico-chemical properties. Due to discrepancies in measurements performed by different methods, confidence in the existing data is low. Data on physico-chemical properties should therefore be used as estimates only (Hertzke *et al.*, 2007). It can be concluded that QA/QC should be carefully considered when generating and interpreting the results of analyses on both toxicological and physicochemical data. Often, uncommon properties of polyfluorinated organic compounds are a result of their structure, which resembles a rigid rod.

The aim of this contribution is a proposal how to estimate the relative toxicity of polyfluorinated compounds. Acute toxicity determination using oligochaeta *Tubifex tubifex* and the relevant mode of approach have been thoroughly described (Tichý & Rucki, 1996; Tichý *et al.*, 2007; 2008). The duration of action of polyfluorinated organic compounds in oligochaeta *Tubifex tubifex* was taken as the basis for this consideration.

Material and methods

Determination of ET_{50}

The perfluorinated acids C_8 (95%), C_9 (97%), C_{10} (98%), C_{11} (95%) and C_{12} (95%) were obtained from Aldrich (Steilheim, Germany). Always fresh oligochaeta *Tubifex tubifex* were purchased in a common shop. Manganese chloride (extra pure grade, Merck) was used as a reference compound (Tichý *et al.*, 2007) to check the quality of the test object.

Solutions of C_8 to C_{12} linear perfluorinated acids were prepared and tested at the concentration of 400 $\mu\text{g/ml}$ (from 0.44×10^{-3} mol/l for C_{12} to 0.58×10^{-3} mol/l for C_8),

which was close to saturated concentrations. The amount of exactly 0.01 g of acid was weighed and dissolved by 15 ml water in a 25 ml plastic volumetric flask and sonicated 10 min at room temperature (22–24 °C). The resulting aqueous acidic solution (pH 3 to 4) was neutralized by titration with a solution of sodium hydroxide (0.1 mol/l) to about pH 7 (checked by pH-meter) and filled up to 25 ml. Neutralization of the solutions was necessary because the low acidity itself could affect the oligochaeta.

ET_n is the effective time when n % of individuals from the population exposed shows a recognizable reaction. In our case, ET_5 , ET_{50} and ET_{95} were determined by counting the number of immobilized *Tubifex tubifex* during the exposure time. Immobilized worms were counted each half hour for five hours of exposure at room temperature (22–24 °C). Deformation of bodies of the oligochaeta was observed: they wrinkled, bled and finally the worms were destroyed and broke up into pieces. Thus immobilization could be noticed only with difficulties. Each solution was tested three times in various seasons and always in a triplicate. The experimental data were processed with special computer software (Logistic Method for Determination of LD_{50} by National Institute of Public Health, Prague) modified for calculation of ET_{50} .

Results

At the concentration of 400 $\mu\text{g/ml}$, aqueous solutions of perfluorinated aliphatic carboxylic acid are not toxic if the acute toxicity is measured in three minutes with *Tubifex tubifex* (Tichý *et al.*, 1996; 2007). The three-minute exposure caused no apparent changes of *Tubifex tubifex*. Thus the solutions of the acids were not acutely toxic for the oligochaeta and the common testing procedure could not be applied.

However, the effect begins to be noticeable after a longer time, in chronic exposure, and the time necessary for induction of observable changes could be taken as a measure of toxicity of the acids. The start of the effect depends on the chain length of the given acid, however with the minimum at C_{11} acid.

The results are summarized in Table 1, which contains values and confidence intervals ($n = 3$, $\alpha = 0.05$) of ET_5 , ET_{50} and ET_{95} for the time necessary to immobilize 5%, 50% and 95%, respectively, of the oligochaeta from the whole population under exposure. The minimum ET_{50} value appeared with perfluoroundecanoic acid (C_{11}) being comparable with perfluorodecanoic acid (C_{10}).

Discussion

We applied an uncommon measure of toxicity, ET_{50} , the time necessary for the onset of destruction of the oligochaeta body. The shorter the ET_{50} , the more toxic is a compound considered. We used this measure because no acute toxicity was observed, a fact that corresponded with the generally accepted opinion that perfluorinated

Table 1. Effective time ET_{50} , ET_5 and ET_{95} (mins.). *Tubifex tubifex* was exposed in neutral aqueous saturated solutions of perfluorinated carboxylic acids (C_8 – C_{12}).

Perfluorinated acid	Effective time (min)		
	ET_5	ET_{50}	ET_{95}
Octanoic $C_7F_{15}COOH$	111 (62–141)	274 (240–347)	675 (471–1687)
Nonanoic $C_8F_{17}COOH$	155 (106–181)	284 (257–339)	519 (405–972)
Decanoic $C_9F_{19}COOH$	23 (12–32)	59 (47–70)	151 (118–233)
Undecanoic $C_{10}F_{21}COOH$	13 (7–18)	32 (25–39)	74 (56–130)
Dodecanoic $C_{11}F_{23}COOH$	13 (7–18)	176 (143–221)	1073 (627–3328)

values in brackets represent confidence intervals ($n = 3$, $\alpha = 0.05$)

carboxylic acids have limited toxicity for aquatic organisms.

Using the ET_{50} measure, the most toxic of the set of the compounds tested, i.e. linear polyfluorinated carboxylic acids, were perfluorodecanoic (C_{10}) and perfluoroundecanoic (C_{11}) acids. No statistically significant differences between the toxic indices exist among C_8 , C_9 and C_{12} acids. The C_{10} and C_{11} acids form a clear minimum, statistically significantly distinguished, and thus considered the most toxic ones in chronic exposures. Naturally, this conclusion applies only for our laboratory conditions in a closed system (on Petri dishes).

The question arises how does $\log P_{(n\text{-octanol-water})}$ of these acids or their salts behave. Unfortunately, even with simpler alcohols and fluorotelomers, it is still an unsolved problem because of the unique behavior of the fluorine substituents in the compounds. As far as the image of perfluorinated acids as rigid rods is true, a bending or a break may exist at C_{10} and C_{11} . We can thus just speculate that there is some non-linearity at the C_{10} to C_{11} carbon chain. The ability to penetrate membranes by this special form of compounds could then be influenced by this fact, although a better penetrating *Tubifex* surface with a rod is more probable than with anything else. The effect in the oligochaeta starts much sooner with C_{10} and C_{11} perfluorinated acids than with shorter or longer carbon chains.

Acknowledgment

We thank the National Institute of Public Health and the Department of Analytical Chemistry, Faculty of Science, Charles University in Prague for allowing to perform this work in their laboratories and to use their libraries.

Sources funding the project: grant of the Ministry of Education of the Czech Republic no. 2B08075, Projects VZ MSM 0021620857 and RP 14/63 MSMT

REFERENCES

- Alloatti D, Giannini G, Cabri W, Lustrati I, Marzi M, Ciacci A, Gallo G, Tinti MO, Marcellini M, Riccioni T, Guglielmi P, Pisano C. (2008). Synthesis and biological activity of fluorinated Combrestatin analogues. *J. Med. Chem.* **51**: 2708–2721.
- Armitage JM, MacLead M, Cousins IT. (2009). Comparative assessment of the global fate and transport pathways of long-chain perfluorocarboxylic acids (PFCAs) and perfluorocarboxalates (PFCs) emitted from direct sources. *Environ. Sci. Technol.* **43**: 5830–5836.
- Herzke D, Schlabach M, Mariussen E, Uggerud H, Heimstad E. (2007). A literature survey on selected chemical compounds. TA-2238/2007. <http://www.sft.no/publikasjoner/2238/ta2238.pdf>.
- Broxterman HJ, Georgopapadakou NH. (2005). Anticancer Therapeutics: "Addictive" multi-targeted drugs, new drug combinations. *Drug Resistance Updates* **8**: 183–197.
- Houde M, Martin JW, Letcher RJ, Solomon KR, Muir DCG. (2006). Biological monitoring of polyfluoroalkyl substances: A review. *Environmental Science & Technology* **40**: 3463–3473.
- Kovarova J, Svobodova Z. (2008). Perfluorinated compounds: occurrence and risk – A review. *Neurotoxicol. Letters* **29**: 599–608.
- Hariharan R, Bray M, Ganim R, Doest T, Goodwin GW, Taegtmeier H. (1995). Fundamental limitations of [^{18}F]2-deoxy-2-fluoro-D-glucose for assessing myocardial glucose uptake. *Circulation* **91**: 2435–2444.
- Tichý M, Rucki M. (1996). An alternative method for determining acute toxicity of chemicals: cessation of movements in *Tubifex tubifex* worms. *Pracov Lék* **48**: 225–230 [in czech].
- Tichý M, Rucki M, Hanzlíková I, Roth Z. (2007). The *Tubifex tubifex* assay for the determination of acute toxicity. *ATLA* **35**: 229–237.
- Tichý M, Rucki M, Hanzlíková I, Roth Z. (2008). Acute toxicity determination by calculation – *Tubifex tubifex* assay and quantitative structure – activity relationships *Environ. Toxicol. Chem.* **27**: 2281–2286.
- Valigurova R. (2008). Thesis: Toxicity determination of salts of perfluorinated organic acids, Charles University in Prague.