

ORIGINAL ARTICLE

Effects of hydrogen peroxide on diazepam and xylazine sedation in chicks

Yaareb J. MOUSA, Fouad K. MOHAMMAD

Department of Physiology, Biochemistry and Pharmacology, College of Veterinary Medicine, University of Mosul, Mosul, Iraq

ITX050412A03 • Received: 24 March 2012 • Revised: 12 December 2012 • Accepted: 15 December 2012

ABSTRACT

Oxidative stress may cause various neuronal dysfunctions and modulate responses to many centrally acting drugs. This study examines the effects of oxidative stress produced by hydrogen peroxide (H_2O_2) on sedation induced by diazepam or xylazine as assessed in 7–14 day-old chicks. Day-old chicks were provided with either plane tap water (control group) or H_2O_2 in tap water as 0.5% v/v drinking solution for two weeks in order to produce oxidative stress. Spectrophotometric methods were used to determine glutathione and malondialdehyde concentrations in plasma and whole brain. Drug-induced sedation in the chicks was assessed by monitoring the occurrence of signs of sedation manifested as drooping of the head, closed eyelids, reduced motility or immotility, decreased distress calls, and recumbency. The latency to onset of sedation and its duration were also recorded. H_2O_2 treatment for two weeks significantly decreased glutathione and increased malondialdehyde concentrations in plasma and whole brain of the chicks on days 7, 10 and 14 as compared with respective age-matched control groups. H_2O_2 decreased the median effective doses of diazepam and xylazine for the induction of sedation in chicks by 46% and 63%, respectively. Injection of diazepam at 2.5, 5 and 10 mg/kg, i.m. or xylazine at 2, 4 and 8 mg/kg, i.m. induced sedation in both control and H_2O_2 -treated chicks in a dose dependent manner, manifested by the above given signs of sedation. H_2O_2 significantly decreased the latency to onset of sedation in chicks treated with diazepam at 5 and 10 mg/kg, increased the duration of sedation and prolonged the total recovery time in comparison with respective non-stressed control chicks. A similar trend occurred with xylazine in the H_2O_2 -treated chicks, though the differences from control counterparts did not attain the statistical significance, except for the recovery time of the lowest dose of the drug. The data suggest that H_2O_2 -induced oxidative stress sensitizes the chicks to the depressant action of the sedatives diazepam and xylazine. Further studies are needed to examine the potential role of oxidative stress in modulating the actions of therapeutic agents on the brain.

KEY WORDS: oxidative stress; sedation; diazepam; xylazine; H_2O_2

Introduction

There is ample evidence indicating that reactive oxygen species are involved in various neuronal disorders and diseases (Aksenova *et al.*, 2005; Kunwar & Priyadarsini, 2011). Oxidative stress (OS) may cause various central nervous system (CNS) dysfunctions (Aksenova *et al.*, 2005; Sayre *et al.*, 2008; Ghosh *et al.*, 2011). Stressful conditions involving the brain may also modulate the response to many centrally acting therapeutic agents (Watanabe *et al.*, 1996; Aksenova *et al.*, 2005). Hydrogen peroxide (H_2O_2) is used experimentally *in vivo* to induce

OS in various laboratory animal species (Desagher *et al.*, 1996; Servitja *et al.*, 2000; Celik & Ozkaya, 2002; Ahmed, 2010; Al-Baggou *et al.*, 2011) and *in vitro* for experimental models of OS in cells and tissues (Frantseva *et al.*, 1998; Aksenova *et al.*, 2005; Gonzalez *et al.*, 2006). H_2O_2 -induced OS was reported to modify the response of rabbits to ketamine-detomidine anesthesia (Wohaieb *et al.*, 1994) and that of rats to pentobarbital anesthesia (Mohammad *et al.*, 1999). A recent study also reported that H_2O_2 potentiated the anticholinesterase poisoning induced by the organophosphate insecticides dichlorvos and diazinon in chicks (Al-Baggou *et al.*, 2011).

The neurotoxic effects of H_2O_2 , through production of free radicals, might be related to lipid peroxidation in the CNS (Piantadosi & Tatro, 1990; Watt *et al.*, 2004) and elevation of intracellular calcium (Hinshaw *et al.*, 1993; Gonzalez *et al.*, 2006). The latter effect could be related to activation of the non-selective cation channel of the cell (Smith *et al.*, 2003). Neurotransmitters, such as

Correspondence address:

Fouad K. Mohammad, BVMS., MS., PhD.

Department of Physiology, Biochemistry and Pharmacology,
College of Veterinary Medicine, University of Mosul,
P. O. Box 11136, Mosul, Iraq.

TEL.: +964-770-160-5334 • E-MAIL: fouadmohammad@yahoo.com

glutamate and GABA, could also be the target of H₂O₂ in the brain (Aksenova *et al.*, 2005; Gonzalez *et al.*, 2006). In this context, the response of the brain to centrally acting therapeutic agents is expected to be modified. Hence, the purpose of the present study was to examine the effects of OS induced by H₂O₂ on diazepam- and xylazine-induced sedation in chicks. Both drugs are centrally acting sedatives in animals, including avian species (Crowell-Davis & Murray, 2006; Rock, 2007).

Materials and methods

Animals

Day-old Cobb broiler chicks of both sexes (body weight 40–90 g) were used. They were maintained in a room with a temperature of 32–35 °C, constant lighting and wood shavings as floor litter. The chicks had free access to drinking water and feed throughout the experiment. All experiments complied with our institutional regulations addressing animal use, and the chicks received proper attention and humane care. The Scientific Committee of the College of Veterinary Medicine at the University of Mosul reviewed and approved the protocol of this study.

Induction of OS

Day-old chicks were either provided with plane tap water (control group) or H₂O₂ (Thomas Baker Chemical Ltd., U.K.) in tap water as 0.5% v/v drinking solution for two weeks in order to produce OS as reported before (Mohammad *et al.*, 1999; Ahmed, 2010). We changed the drinking water and supplied it freshly to the chicks on each day. On treatment days 7, 10 and 14, blood samples were obtained from the chicks (8/group) by jugular vein bleeding into heparinized test tubes (Stevens & Ridgway, 1966). Thereafter, the chicks were euthanized by cervical dislocation to obtain the whole brain. Blood samples were centrifuged at 10 000 rpm for 15 minutes to obtain plasma. Plasma and whole brain samples were kept at –20 °C pending analysis within one week. Spectrophotometric methods were used to determine glutathione concentration in the brain by a modified Ellman method (Ellman, 1959; James *et al.*, 1982) and malondialdehyde concentration was determined by the method of Ohkawa *et al.* (1979). We dissolved diazepam (kindly donated by the State Company for Drugs and Medical Appliances-Ninevah, Iraq) in warm propylene glycol, and further diluted xylazine (2%, Alfasan Co., Holland) in normal saline solution to obtain the desired concentrations of the drugs for injection at a volume of 5 ml/kg body weight given intramuscularly (i.m.).

Effect of H₂O₂ on sedation induced by diazepam and xylazine

The up-and-down method (Dixon, 1980) was used to determine the median effective doses (ED50s) of diazepam and xylazine for the induction of sedation in chicks given tap water (control) or H₂O₂ as described above. The age of the chicks assessed in this experiment

ranged between 7–14 days. The initial dose of diazepam or xylazine was at 10 mg/kg, i.m. After the injection of each drug, we monitored the chicks for the occurrence of sedation manifested as drooping of the head, closed eyelids, reduced motility or immotility, decreased distress calls, and recumbency (Al-Zubaidy & Mohammad, 2005). In another experiment, we monitored the dose-response sedative effects of diazepam and xylazine in the two groups of H₂O₂-stressed and non-stressed chicks. The chicks (8/dose group) were treated i.m. with diazepam at 2.5, 5 and 10 mg/kg or xylazine at 2, 4 and 8 mg/kg. After injection of each drug, the chicks were individually monitored to record the onset of sedation (drooping of the head) and its duration, as well as the total recovery time as described previously (Al-Zubaidy & Mohammad, 2005). Recovery time was the time from the onset of sedation until the chick moved freely. Chicks treated with the vehicle only (propylene glycol or normal saline at 5 ml/kg, i.m.) were also included in all the experiments.

Statistics

The data were statistically analyzed by analysis of variance followed by the least significant difference test (Petrie & Watson, 1999). The level of significance was at $p < 0.05$.

Results

H₂O₂ treatment in drinking water for two weeks significantly decreased glutathione and increased malondialdehyde concentrations in plasma and whole brain of the chicks on days 7, 10 and 14 as compared with respective age-matched control groups (Tables 1 and 2). The ED50s of diazepam and xylazine for the induction of sedation in control chicks, as determined by the up-and-down method, were 6.5 and 4.8 mg/kg, i.m., respectively. In the H₂O₂-treated chicks the sedative ED50s of the drugs decreased by 46% and 63% to 3.5 and 1.8 mg/kg, i.m., respectively.

Injections of diazepam and xylazine induced sedation in the control chicks in a dose dependent manner within 2.6 to 5.1 and 1.6 to 2.5 min, respectively (Tables 3 and 4). The chicks manifested signs of sedation characterized by drooping of the head and wings, closed eyelids, reduced motility or immotility, recumbency and decreased distress calls. The duration of sedation ranged between 55 to 92.3 min for diazepam and between 23.1 to 46.9 min for xylazine, with total recovery times of 82.4 to 115.3 and 27.5 to 65.9 min, respectively (Tables 3 and 4). None of the signs of sedation appeared in the vehicle-treated chicks. H₂O₂ significantly decreased the latency to onset of sedation in chicks treated with diazepam at 5 and 10 mg/kg, increased the duration of sedation and prolonged the total recovery time in comparison with respective non-stressed control chicks (Table 3). We observed a similar trend with xylazine in the H₂O₂-treated chicks, though the differences with control counterparts did not attain the statistical significance, except at the recovery time of the lowest dose of the drug (Table 4).

Table 1. Glutathione concentrations in plasma and whole brain of chicks treated with hydrogen peroxide (0.5%) in drinking water for two weeks.

Groups	Age (days)		
	7	10	14
Plasma (μmol/ml)			
Control	0.213±0.032	0.261±0.022	0.269±0.023
H ₂ O ₂	0.062±0.007*	0.137±0.015 ^a	0.143±0.010 ^a
Whole brain (μmol/g)			
Control	0.477±0.076	0.571±0.039	0.911±0.100 ^{ab}
H ₂ O ₂	0.187±0.025*	0.352±0.036 ^a	0.458±0.048 ^a

Values are mean ± SE for 8 chicks/group
 Hydrogen peroxide was added to drinking water of one-day-old chicks and continued for 14 days.
^{*}Significantly different from the respective control value, $p < 0.05$.
^aSignificantly different from the respective 7-day-old group, $p < 0.05$.
^bSignificantly different from the respective 10-day-old group, $p < 0.05$.

Table 2. Malondialdehyde concentrations in plasma and whole brain of chicks treated with hydrogen peroxide (0.5%) in drinking water for two weeks.

Groups	Age (days)		
	7	10	14
Plasma (nmol/ml)			
Control	3.3±0.59	3.7±0.63	8.5±2.01 ^{ab}
H ₂ O ₂	13.9±0.64*	15.5±3.14*	26.5±2.89 ^{ab}
Whole brain (nmol/g)			
Control	123.9±7.75	138.5±7.61	125.1±6.37
H ₂ O ₂	182.1±4.5*	171.8±12.08*	163.9±7.45*

Values are mean ± SE for 8 chicks/group
 Hydrogen peroxide was added to drinking water of one-day-old chicks and continued for 14 days.
^{*}Significantly different from the respective control value, $p < 0.05$.
^aSignificantly different from the respective 7-day-old group, $p < 0.05$.
^bSignificantly different from the respective 10-day-old group, $p < 0.05$.

Table 3. Sedative effect of diazepam in chicks treated with hydrogen peroxide (0.5%) in drinking water for two weeks.

Diazepam (mg/kg, i.m.)	Duration of sedation (min)	Latency to onset of sedation (min)	Recovery time (min)
Control (tap water)			
2.5	55.0±4.74	5.1±0.55	82.4±3.91
5	74.6±4.44 ^a	4.1±0.77	94.4±6.24
10	92.3±6.47 ^{ab}	2.6±0.38 ^a	115.3±6.64 ^{ab}
Hydrogen peroxide (0.5% in water)			
2.5	44.5±6.32	4.4±0.75	79.3±9.70
5	94.9±6.48 ^a	2.5±0.19 ^a	121.9±4.02 ^a
10	124.9±6.72 ^{ab}	1.3±0.16 ^a	203.5±7.45 ^{ab}

Values are mean ± SE for 8 chicks/group
 Hydrogen peroxide was added to drinking water of one-day-old chicks and continued for 14 days.
^{*}Significantly different from the respective control value, $p < 0.05$.
^aSignificantly different from the respective 2.5 mg/kg dose group of diazepam within the same water treatment regimen, $p < 0.05$.
^bSignificantly different from the respective 5 mg/kg dose group of diazepam within the same water treatment regimen, $p < 0.05$.

Table 4. Sedative effect of xylazine in chicks treated with hydrogen peroxide (0.5%) in drinking water for two weeks.

Xylazine (mg/kg, i.m.)	Duration of sedation (min)	Latency to onset of sedation (min)	Recovery time (min)
Control (tap water)			
2	23.1±1.00	2.5±0.33	27.5±1.39
4	34.1±2.62 ^a	1.9±0.23	45.1±4.70 ^a
8	46.9±4.04 ^{ab}	1.6±0.74 ^a	65.9±8.64 ^{ab}
Hydrogen peroxide (0.5% in water)			
2	27.5±3.05	1.8±0.31	37.1±4.33 [*]
4	39.0±2.54 ^a	1.5±0.19	54.3±4.68 ^a
8	52.1±3.10 ^{ab}	1.3±0.25	75.9±7.8 ^{ab}

Values are mean ± SE for 8 chicks/group
 Hydrogen peroxide was added to drinking water of one-day-old chicks and continued for 14 days.
^{*}Significantly different from the respective control value, $p < 0.05$.
^aSignificantly different from the respective 2 mg/kg dose group of xylazine within the same water treatment regimen, $p < 0.05$.
^bSignificantly different from the respective 4 mg/kg dose group of xylazine within the same water treatment regimen, $p < 0.05$.

Discussion

The decrease in glutathione and increase in malondialdehyde concentrations in plasma and whole brain of the chicks after oral exposure to H₂O₂ suggest that the treatment in drinking water induced OS in the chicks, including their CNS. Glutathione and malondialdehyde are indirect markers of OS in blood and tissues (Janero, 1990; Kohen & Nyska, 2002; Aksenova *et al.*, 2005; Santi *et al.*, 2011). Glutathione is an oxygen radical scavenger, whereas malondialdehyde is the by-product of membrane peroxidation that results in tissue damage and further indicates reduced effectiveness of the mechanisms of antioxidants in protecting the tissue from oxidative damage (Sies, 1999; Karadeniz *et al.*, 2007; Limon-Pacheco *et al.*, 2007). The OS appeared unequivocally on days 7, 10 and 14 of exposure.

These results are in accordance with the reported findings that H₂O₂ oral exposure could be used as a model for the induction of OS in chickens (Ahmed, 2010; Al-Baggou *et al.*, 2011) as well as in rodent species (Desagher *et al.*, 1996; Servitja *et al.*, 2000; Celik & Ozkaya, 2002). The CNS is particularly susceptible to OS-induced neurotoxicity (Sayre *et al.*, 2008). H₂O₂ differentially accumulates in various brain regions (Piantadosi and Tatro, 1990), sets free oxygen radicals that might induce cellular and DNA damage (Gutteridge, 1995; Lee and Jeong, 2007; Xiaoyan *et al.*, 2010; Kunwar & Priyadarsini, 2011). It is also possible that neuronal function is altered as a result of lipid peroxidation, which changes membrane fluidity and damages the protein components of neurons (Kunwar &

Priyadarsini, 2011). Aldehydes and ketones are toxic products of protein oxidation (Kunwar & Priyadarsini, 2011). Furthermore, in vitro experiments showed that oxidative stress caused phospholipid damage in brain slices and astrocytes (Servitja *et al.*, 2000). Oxidative stress as a form of neurotoxic condition (Sayre *et al.*, 2008) was reported to severely damage neurotransmitter signaling systems and modulate enzyme activities, e.g. cholinesterases in the CNS, which might in turn produce an additional burden on neuronal functions (Schallreuter *et al.*, 2004; de Jongh *et al.*, 2007).

Chicks exposed orally to H₂O₂ appeared to be more susceptible to the sedative effects of diazepam and xylazine. This was evident by the decreases in the ED₅₀s of both drugs for the induction of sedation. Further, H₂O₂-stressed chicks showed prolonged sedation induced by the drugs. Diazepam and xylazine are CNS active drugs with different mechanisms of action (Crowell-Davis & Murray, 2006; Rock, 2007). Diazepam action is mediated by potentiation of GABA-ergic inhibition in the CNS, whereas that of xylazine is mediated by activation of alpha₂-adrenoceptors resulting in reduced catecholamine release and turnover (Greene & Thurmon, 1988; Crowell-Davis & Murray, 2006). Corroborating our findings, H₂O₂ treatment in drinking water was also found to sensitize rats and rabbits to the sedative and anesthetic actions of pentobarbital and detomidine-ketamine, respectively (Wohaieb *et al.*, 1994; Mohammad *et al.*, 1999). Other non-chemical stressful conditions, such as restraining and swimming, were reported to alter the pharmacological responses of mice, rats or chicks to diazepam (Marin & Arce, 1996; Kalman *et al.*, 1997; Motohashi *et al.*, 2008). Furthermore, H₂O₂ potentiated the toxicity of organophosphate insecticides acting mainly through a central mechanism in chicks, irrespective of the extent of cholinesterase inhibition (Al-Baggou *et al.*, 2011). In this context, the data of the present study suggest that H₂O₂-induced OS sensitizes the chicks (as a model) to the depressant actions of the sedative drugs diazepam and xylazine. Further studies are needed to get insight into the potential role of OS in modulating the actions of therapeutic agents on the CNS.

Acknowledgements

This report represents a portion of a dissertation submitted by the first author to the University of Mosul, Iraq as partial fulfillment of the requirements for the PhD degree in Veterinary Pharmacology. The study was supported by the College of Veterinary Medicine, University of Mosul, Iraq.

REFERENCES

- Ahmed LI. (2010). *Neurobehavioral and biochemical studies of hydrogen peroxide-induced oxidative stress in chicks*. MSc Thesis. University of Dohuk, Dohuk, Iraq.
- Aksenova MV, Aksenov MY, Mactutus CF, Booze RM. (2005). Cell culture models of oxidative stress and injury in the central nervous system. *Current Neurovascular Res* **2**: 73–89.
- Al-Baggou BK, Naser AS, Mohammad FK. (2011). Hydrogen peroxide potentiates organophosphate toxicosis in chicks. *HVM Bioflux* **3**:142–149.
- Al-Zubaidy MHI, Mohammad FK. (2005). Metoclopramide induced central nervous system depression in the chicken. *BMC Vet Res* **1**: 6–10.
- Celik S, Ozkaya A. (2002). Effects of intraperitoneally administered lipoic acid, vitamin E, and linalool on the level of total lipid and fatty acids in guinea pig brain with oxidative stress induced by H₂O₂. *J Biochem Mol Biol* **35**: 547–552.
- Crowell-Davis S, Murray T. (2006). *Veterinary psychopharmacology*. Blackwell Publishing Professional, Ames, Iowa, USA.
- de Jongh R, Haenen GR, van Koeveeringe GA, Dambros M, De Mey JG, van Kerrebroeck PE. (2007). Oxidative stress reduces the muscarinic receptor function in the urinary bladder. *NeuroUrol Urodyn* **26**: 302–308.
- Desagher S, Glowinski J, Pre'mont J. (1996). Astrocytes protect neurons from hydrogen peroxide toxicity. *J Neurosci* **16**: 2553–2562.
- Dixon WJ. (1980). Efficient analysis of experimental observations. *Ann Rev Pharmacol Toxicol* **20**: 441–462.
- Ellman GL. (1959). Tissue sulfhydryl groups. *Arch Biochem Biophys* **82**: 70–77.
- Frantseva MV, Perez Velazquez JL, Carlen PL. (1998). Changes in membrane and synaptic properties of thalamocortical circuitry caused by hydrogen peroxide. *J Neurophysiol* **80**:1317–1326.
- Ghosh N, Ghosh R, Mandal SC. (2011). Antioxidant protection: A promising therapeutic intervention in neurodegenerative disease. *Free Radic Res* **45**: 888–905.
- Gonzalez A, Granados MP, Pariente JA, Salido GM. (2006). H₂O₂ mobilizes Ca²⁺ from agonist- and thapsigargin-sensitive and insensitive intracellular stores and stimulates glutamate secretion in rat hippocampal astrocytes. *Neurochem Res* **31**:741–750.
- Greene SA, Thurmon JC. (1988). Xylazine - A review of its pharmacology and use in veterinary medicine. *J Vet Pharmacol Therap* **11**: 295–313.
- Gutteridge JMC. (1995). Lipid peroxidation and antioxidants as biomarkers of tissue damage. *Clin Chem* **41**: 1819–1828.
- Hinshaw DB, Miller MT, Omann GM, Beals TF, Hyslop PA. (1993). A cellular model of oxidant-mediated neuronal injury. *Brain Res* **25**:615: 13–26.
- James RC, Goodman DR, Harbison RD. (1982). Hepatic glutathione and hepatotoxicity: changes induced by selected narcotics. *J Pharmacol Exp Therap* **221**: 708–714.
- Janero DR. (1990). Malondialdehyde and thiobarbituric acid-reactivity as diagnostic indices of lipid peroxidation and peroxidative tissue injury. *Free Radic Biol Med* **9**: 515–540.
- Kalman BA, Kim PJ, Cole MA, Chi MS, Spencer RL. (1997). Diazepam attenuation of restraint stress-induced corticosterone levels is enhanced by prior exposure to repeated restraint. *Psychoneuroendocrinology* **22**: 349–360.
- Karadeniz A, Yildirim A, Çelebi F. (2007). Protective effect of panax ginseng against carbon tetrachloride (CCl₄)- induced oxidative brain injury in rats. *Atatürk Üniversitesi Vet Bil Derg* **2**: 117–121.
- Kohen R, Nyska A. (2002). Oxidation of biological systems: oxidative stress phenomena, antioxidants, redox reactions, and methods for their quantification. *Toxicol Pathol* **30**: 620–630.
- Kunwar A, Priyadarsini KI. (2011). Free radicals, oxidative stress and importance of antioxidants in human health. *J Med Allied Sci* **1**: 53–60.
- Lee KJ, Jeong HG. (2007). Protective effect of kahweol and cafestol against hydrogen peroxide-induced oxidative stress and DNA damage. *Toxicol Lett* **173**: 80–87.
- Limon-Pacheco JH, Hernandez NA, Fanjul-Moles ML, Gonshebbat ME. (2007). Glutathione depletion activates mitogen activated protein kinase (MAPK) pathways that display organ-specific responses and brain protection in mice. *Free Radic Biol Med* **43**:1335–1347.
- Marin RH, Arce A. (1996). Benzodiazepine receptors increase induced by stress and maze-learning performance in chick forebrain. *Pharmacol Biochem Behav* **53**: 581–584.
- Mohammad FK, Tawfeek FKh, Hassan AA. (1999). Pentobarbital anesthesia in rats treated with hydrogen peroxide: effect of vitamin E. *Iraqi J Vet Sci* **12**: 203–210.
- Motohashi N, Okamoto Y, Osada M, Yamawaki S. (1993). Acute swim stress increases benzodiazepine receptors, but not GABA_A or GABA_B receptors, in the rat cerebral cortex. *Neurochem Intern* **23**: 327–330.

- Ohkawa H, Ohishi N, Yagi K. (1979). Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal Biochem* **95**: 351–358.
- Petrie A, Watson P. (1999). *Statistics for Veterinary and Animal Science*. Blackwell Science Ltd, Oxford.
- Piantadosi CA, Tatro LG. (1990). Regional H₂O₂ concentration in rat brain after hyperoxic convulsions. *J Appl Physiol* **69**:1761–1766.
- Rock A. (2007). *Veterinary pharmacology. A practical guide for the veterinary nurse*. Butterworth, Heinemann, Elsevier, Edinburgh, U.K.
- Santi A, Menezes C, DuarteMMF, LeitempergerJ, Lópes T, Loro VL. (2011). Oxidative stress biomarkers and acetylcholinesterase activity in human erythrocytes exposed to clomazone (in vitro). *Interdiscip Toxicol* **4**: 149–153.
- Sayre LM, Perry G, Smith MA. (2008). Oxidative stress and neurotoxicity. *Chem Res Toxicol* **21**: 172–188.
- Schallreuter KU, Elwary SMA, Gibbons NCJ, Rokos H, Wooda JM. (2004). Activation/deactivation of acetylcholinesterase by H₂O₂: more evidence for oxidative stress in vitiligo. *Biochem Biophys Res Commun* **315**:502–508.
- Servitja J-M, Masgrau R, Pardo R, Sarri E., Picatoste F. (2000). Effects of oxidative stress on phospholipid signaling in rat cultured astrocytes and brain slices. *J Neurochem* **75**: 788–794.
- Sies H. (1999). Glutathione and its role in cellular functions. *Free Radic Biol Med* **27**: 916–921.
- Smith MA, Herson PS, Lee K, Pinnock RD, Ashford MLJ. (2003). Hydrogen-peroxide induced toxicity of rat striatal neurones involves activation of a non-selective cation channel. *J Physiol* **547**: 417–425.
- Stevens RWC, Ridgway GJ. (1966). A technique for bleeding chickens from the jugular vein. *Poult Sci* **45**: 204–205.
- Watanabe K, Okatani K, Sagara Y. (1996). Potentiating effect of hydrogen peroxide on the serotonin-induced vasocontraction in human umbilical artery. *Acta Obst Gynecol Scand* **75**: 783–789.
- Watt BE, Proudfoot AT, Vale JA. (2004). Hydrogen peroxide poisoning. *Toxicol Rev* **23**:51–57.
- Wohaieb SA, Mohammad FK, Nadir HH. (1994). Effects of hydrogen peroxide-induced oxidative stress on detomidine-ketamine anesthesia in male rabbits. *Iraqi J Vet Sci* **7**:19–23.
- Xiao-yan W, Pei-ying H, Juan D, Jian-zhong Z. (2010). Quercetin in combating H₂O₂ induced early cell apoptosis and mitochondrial damage to normal human keratinocytes. *Chin Med J* **123**: 532–553.